VACCINE INFORMATION SHEET

Polio Vaccine

(IMOVAX® Polio Vaccine)

This vaccine protects against poliomyelitis – a serious infection

- Inactivated polio vaccine (IPV) is given from 2 months of age onwards to protect individuals
 against polio disease. Combination vaccines that contain polio vaccine is routinely given to
 infants and children
- IPV vaccine may be given to children and adults with an incomplete series in order to bring them up-to-date with their immunization
- Proof of vaccination against polio, or a valid exemption, is required by law for all children attending school in Ontario
- The vaccine is also recommended for people who are travelling to countries where risk of polio still exists

Mild side effects may occur, such as:

- Redness, swelling or pain at the injection site are the most common
- Fever, chills, headache and fatigue

People should avoid this immunization if they:

- Have a high fever or serious infection worse than a cold; vaccination should be delayed until
 they are better
- Have ever had a severe allergic (anaphylactic) reaction to a previous dose of this vaccine
- Have an allergy to any vaccine component such as, neomycin, streptomycin, polymyxin B

People should consult their health care provider before vaccination if they:

Have complicated medical issues or previous severe reactions to vaccines

Poliomyelitis (Polio)

- Is a serious disease that can cause nerve damage or paralysis for life
- The virus is spread by contaminated food or water
- Canada has been polio free for the last 20 years but as long as polio exists in other countries, it can be imported to Canada



1-877-464-9675 TTY 1-866-512-6228 york.ca/immunizations

